



# NTP Nonneoplastic Lesion Atlas

## Lung – Amyloid



**Figure Legend:** **Figure 1** Lung - Amyloid in a female B6C3F1/N mouse from a chronic study. There is perivascular and peribronchiolar amyloid deposition (arrows). **Figure 2** Lung - Amyloid in a female B6C3F1/N mouse from a chronic study (higher magnification of Figure 1). There is perivascular and peribronchiolar, acellular, eosinophilic material (arrows). **Figure 3** Lung - Amyloid in a male B6C3F1/N mouse from a chronic study. There is deposition of amyloid in the vascular wall (arrows). **Figure 4** Lung - Amyloid in a male B6C3F1/N mouse from a chronic study. Acellular, eosinophilic material (amyloid) is present in the vessel wall (arrow).

**Comment:** Amyloid (Figure 1, Figure 2, Figure 3, and Figure 4) refers to insoluble, extracellular and/or intracellular proteins that appear as amorphous, eosinophilic, acellular material histologically. It stains



# NTP Nonneoplastic Lesion Atlas

## *Lung – Amyloid*

positively with Congo red stain and has a characteristic apple green birefringence under polarized light. Deposition of amyloid can be focal, multifocal, or diffuse. Involvement of vascular walls is not uncommon, but it may also be seen in peribronchial areas. In severe cases, the pulmonary parenchyma may be effaced. Approximately 90% of the deposits consist of amyloid fibrils formed by the aggregation of misfolded proteins, and 10% of the deposits consist of glycosaminoglycans, apolipoprotein E, and serum amyloid P. Amyloidosis is found as a background change in some strains, such as CD-1 mice. Pulmonary amyloidosis is a spontaneous lesion in aging rodents and may be localized to the lung, or it may be a component of systemic amyloidosis involving spleen, kidney, liver, and other organs. It may also be associated with chronic inflammation. Since an association between amyloid deposition in the lung and xenobiotic administration has not been reported, amyloid deposition is generally considered a background finding. It may, however, be the cause of morbidity or mortality in some animals and can affect the outcome of a study.

**Recommendation:** Because this lesion is biologically important, Lung - Amyloid should be diagnosed and assigned a severity grade whenever present. The location of the amyloid deposits should be indicated in the diagnosis by using a site modifier. Any secondary lesions, such as necrosis or degeneration, should not be diagnosed separately unless warranted by severity but should be described in the pathology narrative.

### **References:**

Buxbaum JN. 2004. The systemic amyloidoses. *Curr Opin Rheumatol* 16:67-75.

Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/14673392>

Frith CH, Manik C. 1991. Incidence, distribution, and morphology of amyloidosis in Charles Rivers CD-1 mice. *Toxicol Pathol* 19:123-127.


Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/1771365>

Higuchi K, Matsumura A, Honma A, Takeshita S, Hashimoto K, Hosokawa M, Yasuhira K, Takeda T. 1983. Systemic senile amyloid in senescence-accelerated mice. A unique fibril protein demonstrated in tissues from various organs by the unlabeled immunoperoxidase method. *Lab Invest* 48:231-240.

Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/6337302>

Homburger F, Russfield AB, Weisburger JH, Lim S, Chak SP, Weisburger EK. 1975. Aging changes in CD-1 HaM/ICR mice reared under standard laboratory conditions. *J Natl Cancer Inst* 55:37-45.

Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/169366>



# NTP Nonneoplastic Lesion Atlas

## *Lung – Amyloid*

### References:

Takeshita S, Hosokawa M, Irino M, Higuchi K, Shimizu K, Yasuhira K, Takeda T. 1982. Spontaneous age-associated amyloidosis in senescence-accelerated mouse (SAM). *Mech Ageing Dev* 20:13-23.  
Abstract: [www.ncbi.nlm.nih.gov/pubmed/7176700](http://www.ncbi.nlm.nih.gov/pubmed/7176700)

Westermarck P, Benson MD, Buxbaum JN, Cohen AS, Frangione B, Ikeda S, Masters CL, Merlini G, Saraiva MJ, Sipe JD. 2007. A primer of amyloid nomenclature. *Amyloid* 14:179-183.  
Abstract: <http://www.ncbi.nlm.nih.gov/pubmed/17701465>

### Authors:

Mark F. Cesta, DVM, PhD, DACVP  
Staff Scientist/NTP Pathologist  
NTP Pathology Group  
National Toxicology Program  
National Institute of Environmental Health Sciences  
Research Triangle Park, NC

Darlene Dixon, DVM, PhD, DACVP  
Group Leader  
Molecular Pathogenesis Group  
National Toxicology Program  
National Institute of Environmental Health Sciences  
Research Triangle Park, NC

Ronald A. Herbert, DVM, PhD  
Group Leader/NTP Pathologist  
Pathology Support Group  
National Toxicology Program  
National Institute of Environmental Health Sciences  
Research Triangle Park, NC

Lauren M. Staska, DVM, PhD, DACVP  
Senior Pathologist  
WIL Research  
Hillsborough, NC